



Pharmacokinetic Metamodel of Morphine and Morphine-6-Glucuronide in Neonates and Adults

Sverrisdóttir, Eva; Foster, David John Richard; Olesen, Anne Estrup; Lund, Trine Meldgaard; Drewes, Asbjørn Mohr; Christrup, Lona Louring; Kreilgaard, Mads; Upton, Richard Neil

Publication date:
2016

Document version
Peer reviewed version

Citation for published version (APA):
Sverrisdóttir, E., Foster, D. J. R., Olesen, A. E., Lund, T. M., Drewes, A. M., Christrup, L. L., ... Upton, R. N. (2016). *Pharmacokinetic Metamodel of Morphine and Morphine-6-Glucuronide in Neonates and Adults*. Poster session presented at PAGE 2016: Population Approach Group Europe, Lissabon, Portugal.



Pharmacokinetic Metamodel of Morphine and Morphine-6-Glucuronide in Neonates and Adults

Eva Sverrisdóttir^{1*}, David John Richard Foster², Anne Estrup Olesen^{1,3}, Trine Meldgaard Lund¹, Asbjørn Mohr Drewes^{3,4}, Lona Lourcing Christrup¹, Mads Kreilgaard¹ & Richard Neil Upton²

¹Department of Drug Design and Pharmacology, Faculty of Health Sciences, University of Copenhagen, Universitetsparken 2, 2100 Copenhagen, Denmark, ²Australian Centre for Pharmacometrics, School of Pharmaceutical and Medical Sciences, City East Campus, North Terrace, University of South Australia, Adelaide SA 5000, Australia, ³Mech-Sense, Department of Gastroenterology & Hepatology, Aalborg University Hospital, Mølleparkvej 4, 9000 Aalborg, Denmark, ⁴Department of Clinical Medicine, Aalborg University, Sdr. Skovvej 11, 9000 Aalborg, Denmark

*E-mail address: eva.sverrisdott@sund.ku.dk

BACKGROUND


Morphine is the gold standard opioid to treat moderate to severe pain. Morphine-6-glucuronide (**M6G**) is an active metabolite and contributes to the effect of morphine [1]. The pharmacokinetics (PK) of morphine and M6G are associated with large inter-individual variability, which makes optimal dose selection challenging, especially in populations such as neonates and patients with renal impairment [2].

[1] Binning, A.R., Przesmycki, K., Sowinski, P., Morrison, L.M., Smith, T.W., Marcus, P., Lees, J.P., Dahan, A., 2011. A randomised controlled trial on the efficacy and side-effect profile (nausea/vomiting/sedation) of morphine-6-glucuronide versus morphine for post-operative pain relief after major abdominal surgery. Eur. J. Pain 15, 402-408.
[2] Somogyi, A.A., Barratt, D.T., Collier, J.K., 2007. Pharmacogenetics of opioids. Clin. Pharmacol. Ther. 81, 429-444.

AIMS

The aims of this study were to develop a **combined morphine and M6G PK metamodel in neonates and adults**, identify covariates that explain some of the large variability, and describe the **formation of M6G** after administration of morphine through different routes.

PHARMACOKINETIC MODEL



METHODS

- Morphine and M6G data from 22 studies, 686 subjects
- Various administration routes
- Healthy volunteers, preterm neonates, and patients with e.g. renal impairment
- NONMEM 7.3.0, PsN, and Pirana
- Sequential modelling of morphine and M6G data

RESULTS

- Two compartment morphine and M6G models
- Two transit compartment absorption for morphine (oral, rectal, and subcutaneous route)
- Systemic and first-pass metabolism of morphine to M6G described
- Several covariates identified including age on morphine clearance, and creatinine clearance on M6G clearance

CONCLUSION

- Some of the large inter-individual variability in morphine and M6G PK was explained with covariates
- Individual morphine and M6G concentrations can be predicted in the shiny application

Parameter	Covariate	Relationship	Equation
Morphine clearance	Postmenstrual age	Sigmoidal	$\frac{PMA^Y}{PMA^Y + TM_{50}^Y} \rightarrow \frac{PMA^{2.31}}{PMA^{2.31} + 113^{2.31} weeks}$
	Age (adults)	Power	$\left(\frac{age}{median\ age}\right)^{AGECOV} \rightarrow \left(\frac{age}{37.5\ years}\right)^{-0.336}$
M6G clearance	Creatinine clearance	Power	$\left(\frac{CR_{CL}}{112.5\ \frac{ml}{min}}\right)^{CRCLCOV} \rightarrow \left(\frac{CR_{CL}}{112.5\ \frac{ml}{min}}\right)^{0.917}$
Morphine and M6G central volume	Venous sampling site	Linear	$(1 + VENC OV) \rightarrow (1 + 0.0936)$
All clearances and volume	Body weight	Power, fixed exponent (0.75 for clearances, 1 for volumes)	$\left(\frac{weight}{70\ kg}\right)^{0.75}$ or $\left(\frac{weight}{70\ kg}\right)^1$

Mean transit time

0.729 h	Oral solution, oral immediate-release tablet, and rectal solution
14.6 h	Oral modified-release tablet
0.310 h	Subcutaneous injection

Bioavailability

14.4%	Oral solution and oral immediate-release tablet
18.8%	Oral modified-release tablet and rectal solution

Systemic metabolism

12.8%	All
-------	-----

First-pass metabolism

20.8%	Oral solution, oral immediate-release tablet, oral modified-release tablet, and rectal solution
-------	---

Link to shiny app: <https://unicph.shinyapps.io/MorphineMetamodel/>

ACKNOWLEDGEMENTS

Data was kindly provided by Albert Dahan, Camilla Staahl, Hanne Villesen, Pernille Ravn, Anne Brokjær, Felix Bochner, Mark Dershitz, Richard Osborne, Magdi Hanna, Ola Dale, Thor Bjelland, Ingolf Meineke, John van den Anker, Brian Anderson.

RESEARCH PARTNERS

